

Learning Enhancement Team

Worksheet: Pythagoras' Theorem


This worksheet has questions about Pythagoras' Theorem which defines the association between the sides of a right-angled triangle. Specifically, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

1. If the hypotenuse of a right-angled triangle is given by c and the other two sides are given by a and b , Pythagoras' Theorem is given by $a^2 + b^2 = c^2$. One problem that can arise when using Pythagoras' Theorem is incorrectly rearranging the equation.


- (i) Rearrange $a^2 + b^2 = c^2$ for a .
- (ii) Rearrange $a^2 + b^2 = c^2$ for b .
- (iii) Rearrange $a^2 + b^2 = c^2$ for c .

Before carrying on, ensure that you have rearranged the equation correctly in each case.


2. Find the missing side in the following triangles. Give your answers to 2 decimal places.




(i)



(ii)




(iii)




(iv)

3. Find the lengths of sides x , y and z in the following triangles. Express your answers as square roots rather than decimal numbers. Triangle (i) is an isosceles triangle and triangle (ii) is equilateral.



(i)



(ii)

4. A Pythagorean Triple is a set of whole number which satisfy Pythagoras' Theorem. The most famous example is $[3, 4, 5]$ so $3^2 + 4^2 = 5^2$. Another example is $[5, 12, 13]$.
- (i) If you double all the numbers in a Pythagorean Triple, do you get another Pythagorean Triple? Check your answer.
- (ii) If you treble all the numbers in a Pythagorean Triple, do you get another Pythagorean Triple? Check your answer.
- (iii) Try some other multiples (both whole numbers, fractions and decimal numbers). Do you still get Pythagorean Triples?
- (iv) Can you show that multiplying each member of a Pythagorean Triple by any number n you still get a Pythagorean Triple? Try to write a sound mathematical argument to show this.